

IVACHENKO, T. I., GRODZENSKIY, D. E.

"The Mineral-Corticoid Function of the Suprarenal Cortex After the Effect of Ionizing Radiation."

Theses of the Proceedings of the Annual Scientific Sessions 23-26 March 1959
(All-Union Institute of Experimental Endocrinology)

From the Radiation Laboratory (Head-Docent D. E. Grodzenskiy) of the All-Union Institute of Experimental Endocrinology (Director--Professor Ye. A. Vasyukova)

G R O - D - E N - K I Y , D . E .

21(b): 17(0) PHASE I: BOOK EXPLANATION 807/2008
International Conference on the Peaceful Uses of Atomic Energy. 24, Geneva, 1955
Booklet available under the Radiobiology I Radiobiology Institute
(Reports of Soviet Scientists, Radiobiology and Radiation Medicine)
Moscow, Izdatvo GIN, 1957. 84 pages. 200 copies. 1000 copies printed.
Soviet Ministry of Health, 1959. 129 p. 2000 copies printed. [Series:
Voprosy kachestvennoy kachestvennoy po mirovuyu 1959. 1000 copies printed.
Study, Vol. 5]

General M.I. A.V. Lebedevskiy, Corresponding Member, USSR Academy of Medical
Sciences; M.I. E.I. Shirokov; Tech. Ed.: Ye.I. Maslov.

PREFACE. This book is intended for physicians, scientists, and engineers
as well as for professors and students at those where radiobiology and
radiation medicine taught.

CONTENTS. This is Volume I of a 6-volume set of reports delivered by Soviet
scientists at the Second International Conference on the Peaceful Uses of
Atomic Energy, held on September 1-13, 1955 in Geneva. Volume I contains
29 reports edited by Candidates of Medical Sciences S.V. Lavitskiy and V.V.
Lebedevskiy. The reports cover problems of the biological effects of ionizing
radiation, the consequences of radiation in small doses, genetic effects
of radiation, the effects of radiation sickness, uses of radioactive isotopes
in medical and biological research, uses of atomic energy for diagnostic
and therapeutic purposes, cell mutagenesis, the use of radioactive isotopes
their intake by plants, and their storage in plants and foodstuffs.
References accompany each report.

Reports of Soviet Scientists (Cont.)

- 160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

GRODZENSKIY, D.E.

Radiation effects and the endocrine system; survey of experimental
data. Med. rad. 4 no.5:77-83 My '59. (MIRA 12:7)

1. Iz Vsesoyuznogo instituta eksperimental'noy endokrinologii.
(RADIATION, eff.
role of endocrine system, review (Rus))
(ENDOCRINE GLANDS, physiol.
role in eff. of radiations, review (Rus))

GRODZENSKIY, D.M.; KROTKOV, P.G.

Review of an article "Six cases of acute radiation sickness in victims of a nuclear reactor accident in Yugoslavia" by H.Jammet and others. Med.rad. 4 no.9:83-90 S '59. (MIRA 12:11)
(RADIATION SICKNESS) (JAMMET, H.)

GRODZENSKIY, D.E.; KROTKOV, F.G.

Review of "Homologous marrow grafts and transfusions in humans accidentally subject to high irradiation" by G. Mathé and others, and "Quantitative determination of the source of erythrocytes in four persons irradiated with large doses and treated by marrow injections" by C. Salmon, Med.rad.4 no.10:85-92 0 '59.

(MIRA 13:2)

(RADIATION SICKNESS) (MARROW--TRANSPLANTATION) (MATHÉ, G.)
(SALMON, C.)

TRET'YAKOVA, K.A.; GRODZENSKIY, D.E.

Biosynthesis of cholesterol and fatty acids in the liver and adrenals
of rats exposed to the effect of ionizing radiations. Vop.med.khim.
5 no.5:362-366 S-O '59. (MIRA 13:2)

1. Radiation Laboratory, the All-Union Institute for Experimental
Endocrinology, Moscow.

(CHOLESTEROL metab.)

(FATTY ACIDS metab.)

(ADRENAL GLANDS radiation eff.)

(LIVER radiation eff.)

TOLKACHVSKAYA, Nadezhda Filippovna; GRODZENSKIY, D.E., red.; BUL'DYAYEV,
H.A., tekhn.red.

[Development of metabolic process in children in the first year
of life] Razvitie protsessov obmena u detei pervogo goda zhizni.
Izd.2. Moskva, Gos.izd-vo med.lit-ry Medgiz, 1960. 256 p.

(MIRA 13:11)

(METABOLISM)

(INFANTS)

VASYUKOVA, Ye.A., pref., red.; GRODZENSKIY, D.E., red.; ZUYEVA, N.K.,
tekhn. red.

[Present-day problems in endocrinology] Sovremennyye voprosy
endokrinologii. Moskva, Medgiz, 1960. 282 p. (MIRA 14:10)
(ENDOCRINOLOGY)

KOZLOVA, A.V., prof., otv.red.; TROITSKIY, V.L., red.; KURLYANDSKAYA,
E.B., red.; BELOUSOV, A.P., red.; IVANITSKIY, A.P., red.;
GRODZENSKIY, D.E., red.izd-va; ASTAF'YEVA, G.A., tekhn.red.

[Medical radiology] Meditsinskaya radiologiya. Moskva, Izd-vo
Akad.nauk SSSR, 1960. 400 p. (MIRA 13:4)

1. Vsesoyuznaya nauchno-tekhnicheskaya konferentsiya po primene-
niyu radioaktivnykh i stabil'nykh izotopov i izlucheniye v narodnom
khozyaystve i nauke, Moscow, 1957. 2. Ministerstvo zdravookhraneniya
SSSR i Institut rentgenologii i radiologii RSFSR, Moskva (for Kozlova).
 3. Institut gigiyeny truda i profzabolevaniy Akademii meditsinskikh
nauk SSSR (for Kurl'yanskaya).
- (BIOLOGY, MEDICAL)

TRET'YAKOVA, K.A.; GRODZENSKIY, D.E.

Effect of thyreiodine and thyroidectomy on the rate of synthesis of cholesterol and fatty acids in rats under the influence of radiation. Vop. med. khim. 6 no. 6:611-614 N-D '60. (MIRA 14:4)

1. Radiation Laboratory of the All-Union Institute of Experimental Endocrinology, Moscow.

(THYROID GLAND) (CHOLESTEROL) (FATTY ACIDS)

(RADIATION SICKNESS)

TRET'YAKOVA, K.A.; GRODZENSKIY, D.E.

The rate of cholesterol and fatty acid synthesis in the adrenals, testes, and liver of young and old rats under normal conditions after irradiation. Biokhimiia 25 no. 3:399-403 My-Je '60.

(MIRA 14:4)

1. Radiatsionnaya laboratoriya Vsesoyuznogo instituta eksperimental'noy endokrinologii, Moskva.

(CHOLESTEROL METABOLISM) (FATTY ACID METABOLISM) (AGING)
(RADIATION—PHYSIOLOGICAL EFFECT)

GRODZENSKIY, D.Ye. (Moskva)

New data on the effect of ionizing radiations on the endocrine
system. Probl.endok.i gorm. 7 no.4:26-32 '61. (MIRA 14:8)
(ENDOCRINE GLANDS) (RADIATION--PHYSIOLOGICAL EFFECT)

GRODZENSKIY, D. E., TRETYAKOVA, K. A., (USSR)

"Effect of Hormonal Factors on the Rate of Synthesis
of Cholesterol in Normal and Irradiated Rats."

Report presented at the 5th Int'l. Biochemistry Congress,
Moscow, 10-16 Aug 1961.

GRODZENSKIY, David Emmanuilovich; ALYAB'YEV, A.F., red.; VLASOVA, N.V., tekhn.
red.

[Radiobiology; biological effect of ionizing radiations] Radiobiologiya;
biologicheskoe deistvie ioniziruiushchikh izluchenii. Moskva, Gos. izd-
vo lit-ry v oblasti atomnoi nauki i tekhniki. 1961. 132 p.

(MIRA 14:8)

(Radiobiology)

GRODZENSKIY, D.

"Problems in the pathogenesis, experimental treatment, and prevention of radiation sickness." Reviewed by D.Grodzenskii.
Pat.fiziol. i eksp. terap. 5 no.3:94-96 My-Je '61. (MIRA 14:6)
(RADIATION SICKNESS)

GRODZENSKIY, D.E.; IVANENKO, T.I.

Changes in the metabolism of electrolytes in rats following
the action of ionizing radiation. Biul. eksp. biol. i med.
51 no.6:62-65 Je '61. (MIPA 15:6)

1. Iz radiatsionnoy laboratorii (zav. - dotsent D.E. Grodzenskiy)
Vsesoyuznogo instituta eksperimental'noy endokrinologii (dir. -
prof. Ye.A. Vasyukova), Moskva. Predstavlena deystvitel'nym
chlenom AMN SSSR F.G. Krotkovym.
(SODIUM METABOLISM) (POTASSIUM METABOLISM)
(X RAYS—PHYSIOLOGICAL EFFECTS)

GRUDZENSKIY, D.E.; KOKOSOV, L.V., red.; VLASOVA, N.A., tekhr. red.

[Isotopes in medicine] Izotopy v meditsine. Moskva, Gos.
izd-vo lit-ry v oblasti atomnoi nauki i tekhniki, 1962. 138 p.
(MIRA 15:3)

(ISOTOPES—THERAPEUTIC USE)

RAYSKINA, Mina Yevgen'yevna; GRODZENSKIY, D.E., red.; BEL'CHIKOVA,
Yu.S., tekhn. red.

[Biochemistry of the nervous regulation of the heart] Biokhi-
mii nervnoi reguliatsii serdtsa. Moskva, Medgiz, 1962. 318 p.
(MIRA 15:9)

(NERVES, CARDIAC)

GRODZENSKIY, David Emmanuilovich; NIKITINA, T.K., red.; MAZEL', Ye.I.,
Lekhn.red.

[Radiobiology; biological effect of ionizing radiations]
Radiobiologiya; biologicheskoe deistvie ioniziruiushchikh izluchenii.
Moskva, Gosatomizdat, 1963. 198 p. (MIRA 16:12)
(RADIOBIOLOGY) (RADIATION—PHYSIOLOGICAL EFFECT)

VASYUKOVA, Ye.A., prof., red.; GRODZENSKIY, D.E., red.; KOKIN,
N.M., tekhn. red.

[Contemporary problems of endocrinology] Sovremennye voprosy
endokrinologii. Moskva, Medgiz. No.2. 1963. 309 p.
(MIRA 16:5)

(ENDOCRINOLOGY)

GRODZENSKIY, D.E.; ZAMYCHKINA, K.S.

Use of the isotope method for the study of absorption from the
digestive tract. Med.rad. no.1:71-76'63. (MIRA 16:10)
(ALIMENTARY CANAL) (ABSORPTION (PHYSIOLOGY)
(TRACERS (BIOLOGY))

BAGRAMYAN, E.P.; GRODZENSKIY, D.E. (Moskva)

Action of X rays and hormones on the penetration of labeled albumin into an inflammatory focus. Pat. fiziol. i eksp. terap. 7 no.6:27-31 N-D '63. (MIRA 17:7)

1. Iz radiatsionnoy laboratorii (zav. - dotsent D.E. Grodzenskiy) Vsesoyuznogo instituta eksperimental'noy endokrinologii.

GROFZENSKIY, D.E.; GORIZONTOV, P.D.; VOROB'YEV, V.I.; ...
FEDOROVA, T.A.; PAVLOVA, M.N.; GABUNIYA, B.I.

Second International Congress on Radiation Research in England,
Aug. 5-11, 1962. Med. rad. 8 no.3:83-92 Mr '63. (MIRA 17:9)

GRODZENSKIY, D.E.

Fifty years of the isotope method. Vop. med. khim. 9 no.5:
540-546 S-O '63. (MIRA 17:1)

SVIRIDOV, N.K.; FATEYEVA, M.N.; GHODZINSKIY, D.E.

Reviews and bibliography. Med. rad. 9 no.11:71-78 N '64.
(MIRA 18:9)

OSIBZHENSKIY, D.E.; IVANKO, T.I.; BAGDASARYAN, V.S.; ALIKHANOVA, I.V.

Biosynthesis of corticosteroids in adrenal tissues in irradiated hypophysectomized rats and electrolyte metabolism. *Izvest. endok. i gorm. 11 no.5:77-81 S-O '65.* (MIRA 1965)

1. Vsesoyuznyy institut eksperimental'noy endokrinologii, Moskva.
Submitted October 20, 1964.

GRODZENSKIY, G.V., inzh.

Initial conditions for calculating allowances for worn hobs
used for cutting cycloidal wheels. Vsesoi tekhn. i inzh.
mashinostr.; meshvuz.sbor. no.3:93-101 '61. (MIRA 14:8)
(Gear-cutting machines)
(Tolerance (Engineering))

1.100

24762

S/119/61/000/007/006/008
D247/D306

AUTHOR: Grodzenskiy, G.V.

TITLE: Small module hard-alloy gear cutters

PERIODICAL: Priborostroyeniye, no. 7, 1961, 21 - 23

TEXT: This article describes a method of making the diamond wheels used in the backing off operation of gear cutters, a method of finding its profile. A new method of manufacture of hard-alloy worm holes and cutting discs suitable for milling pinions and wheels of 0.05 - 0.3 mm module is being widely used at the Pervyy Moskovskiy chasovoy zavod (First Moscow Watch Factory). The milling cutters are made of the hard alloy BK6M - (VK6M). The backing off of cutting discs is carried out on the relieving lathe C-43, - (S-43) and of worm holes on the machine 1810. In both cases the cutter holder is replaced by a high-speed pneumatically driven shaft. A backing off diamond wheel, having a profile corresponding to the profile of the cutter, is fastened to this shaft. The diameter of the wheel is 10 - 20 mm. The

Card 1/4

24762

Small module hard-alloy...

S/119/61/000/007/006/008
D247/D306

speed of revolution is 50,000 rev/min. The backing off is carried out in 2 - 3 revolutions. During the first revolution $2/3$ of the over-measure is removed. On passing from the backing off of the first tooth to the second tooth the speed is increased. The correct choice of the diamond wheel diameter is directed by its correct speed of cutting and durability which also depends on the area of the working surface. On the other hand the diameter must be small enough to give a correct dimension of the backed off part of the tooth ($\angle \theta$) in Fig. 2. The face line of tooth and the path of the corresponding movement of the center of the diamond wheel are equidistant. The formula for the relationship between the maximum value of θ corresponding to the point M and the diameter of the diamond wheel D_k is obtained from the condition that at the point of contact M of the circle with the line KL, this circle must not enter the body of the cutter at the next tooth. If the backing off line KL is an Archimedes spiral, the relationship between D_k and the parameter of the milling cutter

Card 2/4

21.762

Small module hard-alloy...

S/119/61/000/007/006/008
D247/D306

can be expressed by the following practical formulae: (1), (2), (3), (4),

$$D_s = \frac{2 p_M r \cos \beta - p_M^2 - r^2}{p_M \cos \alpha - r \cos (\beta - \alpha)};$$

$$\operatorname{ctg} \alpha = \operatorname{ctg} \alpha_1 - \theta;$$

$$p_M = r(1 - \theta \operatorname{tg} \alpha_1);$$

$$\beta = \frac{2\pi}{z} - \theta;$$

where z is the number of teeth of cutter. The choice of the diameter of the diamond wheel is determined by the limits of the

Card 3/ 4

Small module hard-alloy...

24762
S/119/61/000/007/006/008
D247/D306

mini-mum value of 0 and D_k . The height of the wheel profile measured as a magnitude $\frac{D_k - D_{k'}}{2}$ for a number of cutters is

smaller than the height of the tooth profile in its radial section by 0.01 - 0.04 mm. With a sufficient approximation for all practical purposes the author explains briefly how to find the profile of the diamond wheel. Durability of hard-alloy module milling cutters is 30 or more times higher than that of the steel milling cutters. There are 4 figures.




Fig. 2

Card 4/4

GRODZENSKIJ - G.V.
GRODZENSKIJ, G.V. [Grodzenskiy, G.V.]

Hard metal milling cutters for small-sized gearings. Jemna
mech opt 8 no.6:186-187 Je '63.

SHNEYZERSON, M.B.; GRODZENSKIY, V.A.

One way of interpreting data of the method of reflected waves with
seismographs arranged in groups on large bases. Razved. i prom.
geofiz. no.28:16-21 '59. (MIRA 13:1)
(Prospecting--Geophysical methods)

GRODZENSKIY, V.A.; LEV, I.S.

Installation for combined grouping of PKG-2 seismographs.

Razved.i prom.geofiz. no.4346-51 '62.

(MIRA 15:8)

(Seismometry--Equipment and supplies)

VOYUTSKIY, V.S.; GRODZENSKIY, V.A.

Interference rejection of asynchronous accumulation. Geol. i
geofiz. no.11:90-93 '64. (MIRA 18:4)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut geofizicheskikh
metodov razvedki, Moskva.

... ..

... .. of excitation of elastic
... .. (XRA 18:9)

ACC NR: AT6020745

SOURCE CODE: UR/2552/65/000/046/0021/0033

AUTHOR: Grodzenskiy, V. A.; Beklemishev, A. B.; Kozlova, V. G.

ORG: none

TITLE: Certain findings on the use of the asynchronous accumulation method in seismic prospecting

SOURCE: Moscow. Vsesoyuznyy nauchno-issledovatel'skiy institut geofizicheskikh metodov razvedki. Prikladnaya geofizika, no. 46, 1965, 21-33

TOPIC TAGS: seismic prospecting, hodograph, signal correlation

ABSTRACT: The paper discusses the first application on land of this method which has been used for prospecting at sea since 1959. The work was done in southwest Turkmenistan and was intended to test the applicability of the method under conditions typical of the Transcaspian region. The method was used in two variants. In the first, the signal is recorded in both correlating channels at the same time. In the other, there is a certain time lag between the recordings in the correlating channels. Instrumentation and procedures are briefly discussed. An innovation introduced in the procedure was to do all the shooting at permanent sites and to move the recorders around. The quantities of explosives used and the conditions of shots are tabulated. The quantities of explosives varied from 3 to 4000 kg and the distances between the shot and

Cord 1/2

ACC NR: AT6020745

the receivers varied from 8 to 322 km. In comparison with other seismic methods, the saving in explosives was from 25 to 33%. Traces of waves recorded by this method were sharper and wave velocities were somewhat greater. The criterion for identification of the useful signal is the periodicity of the function of mutual correlation. The results of this experimental work, which had to be abridged because of the climatic conditions, were encouraging. The authors conclude that the method is promising. Orig. art. has: 6 figures.

SUB CODE: 08/ SUBM DATE: none/ ORIG REF: 005

Card 2/2

ACC NR: AT7002054

SOURCE CODE: UR/2552/66/000/047/0029/0041

AUTHORS: Grodzenskiy, V. A.; Lev, I. S.; Slutskovskiy, A. I.

ORG: none

TITLE: The problem of selective properties and sensitivity in groups of low-frequency seismic receivers connected in parallel or in series

SOURCE: Moscow. Vsesoyuznyy nauchno-issledovatel'skiy institut geofizicheskikh metodov razvedki. Prikladnaya geofizika, no. 47, 1966, 29-41

TOPIC TAGS: seismologic instrument, ~~seismic detection~~, seismologic station, seismograph

ABSTRACT: Grouping of seismic detectors has been successfully used to increase their effective sensitivity. Sometimes the grouping is used in order to increase the sensitivity of the seismorecording channel; in that case the seismic detectors are connected in series. However, frequently the desired increase in the sensitivity is not achieved because the true relationships between the output impedance of a group of seismic detectors, the resistance of connecting wires, and the input impedance of amplifiers, i.e., the transient characteristics, are not taken into account. A transient characteristic in this case is defined as the ratio of the frequency characteristic of a seismic detector, coupled to the input of an amplifier, to the frequency characteristic of the same seismic detector under no-load conditions.

Card 1/2

UDC: 550.83(061.6)

ACC NR: AT7002654

The problem of stationary harmonic oscillations in an equivalent "seismic detector - amplifier input" circuit is considered. It is shown that when n identical seismic detectors are connected either in parallel or in series, the phase shift (φ) remains constant; however, the absolute output impedance is decreased n times when the detectors are connected in parallel, and is increased n times when they are connected in series. The absolute impedance and the phase shift were determined experimentally for seismic detectors of the SPEN-1, NS-3, and SP-15 types. It was found that for all three detector types, the absolute output impedance had its largest value at the frequency of the first electromechanical resonance (ω_1); at the same frequency $\varphi = 0$. At frequencies $\omega > 2\omega_1$, the absolute output impedance of all the three detectors varied much less than at $\omega < \omega_1$ where it sharply decreased with a decrease in the frequency. On the basis of the experimental study of the characteristics of NS-3 type detectors, it was found that it is most suitable to connect them in series, thus excluding the resonance conditions of operation. However, when grouping the seismic detectors of the SPEN-1 type, it is most practical to connect them in parallel when they are coupled to seismic stations of low input impedance, and in series when they are coupled to seismic stations of high input impedance. Orig. art. has: 18 formulas, 8 figures, and 2 tables.

SUB CODE: 08 / SUBM DATE: none/ ORIG REF: 002/ ATD PRESS: 5113

Card 2/2

L 40317-66 31(1) 68

ACC NR: AP6005348 SOURCE CODE: UR/0413/66/000/001/0092/0092 32
B

INVENTOR: Voyutskiy, V. S.; Vishnyakov, Ye. P.; Shnirson, M. B.; Levi, I. S.;
Grodzenskiy, V. A.; Tabakov, A. P.

ORG: none

TITLE: Method of recording weak explosions and earthquakes. Class
42, No. 177640

SOURCE: Izobreteniya, promyshlennyye obraztsy, tovarnyye znaki, no. 1,
1966, 92

TOPIC TAGS: earthquake, ~~earthquake recording~~, seismic ~~vibration wave~~,
~~correlation function~~, explosion, ~~explosion recording~~ *seismology*

ABSTRACT: An Author Certificate has been issued for a method of recording
weak explosions and earthquakes based on determination of the interrela-
tion function of seismic vibrations. To improve the quality and reliability
of measurements, the values of the function obtained for a number of
receiving points arranged along the profile are summed up with the vary-
ing time shifts corresponding to those predetermined by the location of
the receiving points along the profile. [LD]

SUB CODE: 08/ SUBM DATE: 29Jan63/

Card 1/111LP

UDC: 550.341

GRUZENKO, V. I., assistant

Studying the formation of fresh-water lenses in coastal zone channels, taking into account the difference in the deposition of fresh and salt waters. Izv. vyznauk. i tekhn. i inzh. S. no. 11:94-98 II '65. (N. 11:94-98)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut gidrogeologii i inzhenernoy geologii.

GRODZENSKIY, Ye.D., inzh., red.; PRVZNER, A.S., red. izd-va; TOKER, A.M.,
tekh. red.

[Manual of consolidated indices of the cost of planning and research]
Spravochnik ukрупnennykh pokazatelei stoimosti proektnykh i issledovatel'skikh rabot. Vvoditsia v deistvie s 1 ianvaria 1958 g. Pt.15.
[Enterprises of the building materials industry] Predpriatia promyshlennosti stroitel'nykh materialov. Moskva, Gos. izd-vo lit-ry po stroit. i arkhitekt. 1957. 92 p. (MIRA 11:8)

1. Russia (1923-
stroitel'stva.

U.S.S.R.) Gosudarstvennyy komitet po delam

(Building materials industry)

FIATO, Stanislaw; GRODZICKA-KROLAK, Hanna; MALCZEWSKI, Bohdan, asystent
techniczny Helena Szymanska

Isolation and typing poliomyelitis virus strains during the 1956
epidemic. Med. dosw. mikrob. 11 no.1:31-38 1959.

1. Z Zakladu Wirusologii PZH - Warszawa.

(POLIOMYELITIS VIRUS,

isolation & typing during epidemic (Pol))

GRODZICKI, Andrzej

Explanation of the sources of trace gold deposits in the
Iagnickie Pole--Nikolajowice--Wadroze Wielkie region.
Mszechswiat no. 1:21-22 Ja '64.

L 41815-66 EWP(j) RM
 ACC NO: AP6031689 (N) SOURCE CODE: PO/0099/66/040/003/0373/0380
 AUTHOR: Swinarski, Antoni; Grodzicki, Antoni 26
 ORG: Department of Inorganic Chemistry, M. Copernicus University, Torun (Katedra B
 Chemii Nieorganicznej Uniwersytetu M. Kopernika)
 TITLE: Determination of the formation constants of chloronitrite complexes of cadmium 1
 by the polarographic method
 SOURCE: Roczniki chemii-annales societatis chimicae polonorum, v. 40, no. 3, 1966,
 373-380
 TOPIC TAGS: intermolecular complex, polarographic analysis
 ABSTRACT: The composition and formation constants of cadmium chlorinitrite complexes
 were determined by the polarographic method. The disproportionation constants of
 the complexes were calculated from the formation constants obtained. Orig. art.
 has: 3 figures, 4 formulas and 3 tables. [Based on authors' Eng. abstr.]
 [JPRS: 36,002]
 SUB CODE: 07 / SUBM DATE: 20May65 / ORIG REF: 001 / SOV REF: 002
 OTH REF: 009

Card 1/1 of

0217 0251

GRODZICKI, J., mgr.; RUDOWSKI, S., mgr

Stereobasis in cartographic-geological research and work. Przegl geod
34 no.9:399 S '62.

1. Zaklad Kartowania Geologicznego, Wydzial Geologii, Uniwersytet,
Warszawa.

GRONICKI, H.

The question of leveling of the 3d and 4th degrees. p. 36.
ASTA PENSICA POLONICA. Warszawa. Vol 12, No. 7, July 1956.

East European Accessions List (EEAL) Library of Congress
Vol. 5, No. 11, August 1956.

GRODZICKI, Zbigniew.

Synthesis of glutamic acid from human hair. Acta Poloniae pharm.
11 no.3:183-187 1954.

1. Z Zakladu Chemii Organicznej Akademii Medycznej w Lublinie.
Kierownik: prof. dr Stanislaw Rolski. Obecni: prof. mgr Andrzej
Szuchnik.

(GLUTAMATES, preparation of,
from human hair)

(HAIR,
preparation of glutamic acid from human hair)

BLITSHTAYN, I.I., kandidat biologicheskikh nauk; MOLDAVSKAYA, V.D., professor;
RODKIN, S.V., dotsent; CHERNYAVSKAYA, F.P., kandidat meditsinskikh nauk;
LEVITAN, R.B.; GRODZINSKAYA, A.I.; OSTROMUKHOVA, B.L.

The role of *Leishmania* and *hymenolepis nana* in dysentery of young
children. Sov.med.21 no.3:22-26 Apr '57. (MLRA 10:7)

1. Iz Ukrainского instituta malyarii i meditsinskoy parazitologii imeni
prof. V.Ya.Rubashkina (dir. I.A.Demchenko), Khar'kovskogo instituta
okhrany materinstva i detstva (dir. - kandidat meditsinskikh nauk A.I.
Kornikova), detskoy bol'nitsy No.24 (glavnyy vrach L.M.Poyarkova) i
detskikh yasley No.81 (glavnyy vrach B.L.Ostromukhova) Khar'kov.

(DYSENTERY, BACILLARY, in inf. and child
in giardiasis & tapeworm infection, ther.)

(GIARDIASIS, in inf. and child
in bacillary dysentery, with tapeworm infect., ther.)

(TAPESWORM INFECTION, in inf. and child
in bacillary dysentery, with giardiasis, ther.)

GRODZINSKAYA, G. S., CAND BIO SCI, A ^{peculiarities} ~~STUDY~~ OF THE ~~CHARACTERISTICS~~ OF THE GROWTH AND DEVELOPMENT OF THE PROGENY OF ~~THE~~ BIOLOGICAL GROUPS OF SUGAR BEETS. KIEV, 1961.
(ACAD SCI UKSSR. DEPT ^{of} BIO SCI). (KL, 2-61, 204).

-76-

GRODZINSKAYA, K. P.: Master Biol Sci (diss) -- "The participation of soil microflora in improving the conditions of manganese nutrition of plants". Kiev, 1959. 16 pp (Acad Sci Ukr SSR, Dept of Biol Sci), 150 copies (KL, No 16, 1959, 107)

ACCESSION NR: AR4027225

8/0299/64/000/002/B044/B044

SOURCE: RZh. Biologiya, Abs. 2B308

AUTHOR: Grodzinskaya, K. P.

TITLE: Transformation of manganese in the soils of the Ukraine under the influence of bacteria

CITED SOURCE: Sb. Primeneniye mikroelementov, polimerov i radioakt. izotopov v s. kh. Vy* p 1. Kiyev, Ukr. akad. s.-kh. nauk, 1962, 145-148

TOPIC TAGS: soil bacterium, manganese, bacterial metabolism, plant physiology, mineral nutrition, soil manganese, rhizosphere

TRANSLATION: The author studied the intensity of the bacterial metabolism of manganese in four types of soil from the Ukraine: soddy podzolized soil, grey podzolized soil, common chernozem and southern chernozem. Samples of soil were taken from the rhizosphere of corn after the appearance of sprouts, during flowering and during harvest time. The total number of microorganisms, the fungi and actinomycetes, the ability of the soil to oxidize manganese, the total and free manganese content, and the pH and soil humidity were determined. It was

1/2

Card

ACCESSION NR: AR4027225

found that there are marked fluctuations in the content of free manganese during growth, due to changes in the ratio of its valence states, which depend on the oxidation and reduction of manganese by various species of microorganisms and their combined activity under the conditions of the rhizosphere of agricultural crops. The transformation of manganese by bacteria was established for all of the soils studied. A. Lambina

DATE ACQ: 14Feb64

SUB CODE: LS

ENCL: 00

2/2

Card

CA

STROZINSKI, J

17

New synthetic estrogens. Jurek-Kluc. *Appl. Zyg. J.*
11, 2 (1965). A review with 28 references. R. H. S.
The complete synthesis of estrogenic hormones. Jurek-Kluc.
Stroinski. *Bull. Acad. Sci. (Chem. 4, 113-21 (1959)).*—A review.
Adam Spawnyński

GRODZINSKA, Krystyna

Meadow and field associations of the Gubalowka Elevation,
Polish Western Carpathians. *Fragm florist* 7 no.2:357-416
'61.

1. Institute of Botany, Polish Academy of Sciences, Krakow.

GRODZINSKA-ZACHWIEJA, Z.; KAHL, W.; PASEK, W.;

Bacteriostatic action of chicory (*Cichorium intybus* L).
Bul Ac Pol biol 10 no.12:513-517 '62.

1. Department of Organic Chemistry, School of Medicine,
Krakow. Presented by J. Heller.

GRODZINSKI, L.Z.

P.O.L.

Contractions of isolated heart of European Glass eel *Anguilla anguilla*. L. Z. Grodzinski (*Bull. Acad. Polon. Sci.*, 1954, 2, 19-22). ~~Heart of European Glass eels~~ in Tyrode's solution modified for cold blooded animals, and with the addition of cell liver homogenate, pulsated within a temp. range of 2.5-40°. The rate increased with the temp. up to 25° but decreased above 30° and temp. above 39° had a lethal effect on the isolated hearts although some which had ceased to pulsate at a lethal temp., began again when the temp. was reduced to room temp. When the sinus venosus was cut off, the atrial pacemaker acted efficiently without the sinus pacemaker and dominated and controlled the ventricular pacemaker, but pulsations were at a lower rate. Partial severance of the tissue conducting the contractions from the auricle to the ventricle lowered the frequency of the ventricular pulsations. When the ventricle was completely severed from the auricle, the main ventricular pacemaker was unable to induce pulsations of its own part of the heart, but, under certain circumstances, contractions were initiated by an auxillary ventricular pacemaker situated in the ventricular dorsal wall near the base of the arterial conus. In the isolated sinus venosus, contractions occurred in a few instances but at a higher rate than in the corresponding auricle and ventricle.

A. ACKROYD.

GRODZINSKI, M. Z.

Susceptibility of the heart in sea trout embryos *Salmo trutta*, L. to small changes of temperature. M. Z. Grodzinski (*C. R. Nat. Acad. Sci., Paris*, 1960, No. 4-8, 17).—The embryos were examined at 5 different growth stages and tested within the range +8° to +18°. All reacted to temp. rises by increasing the heart rate from 4–7 contractions per minute. The max difference of temp to which the majority responded was 0.2°. D S PARWORTH

ASS-SLA METALLURGICAL LITERATURE CLASSIFICATION
140000 STIOLCER
140000 HIF ONE GUE
REVISIONS
1124 8747V
011111 200 151

DIAMOND TOOL PATENTS: II, DIAMOND ABRASIVE WHEELS.
 Edited by P. Grodzinski. Industrial Diamond Information
 Bureau, Industrial Distributors, Ltd., London E.C.1,
 Aug., 1948. 52 pp. Price 10s. — A survey on the use of
 diamonds as abrasives is presented. In particular, the
 type of bonds used in abrasive wheels is discussed. About
 400 abstracts of British, American, German, Russian, and
 Canadian patents are given. These patents also pertain to
 abrasive materials other than diamonds.

GRODZINSKI, P.

4

Developments in hardness testing. P. Grodzinski *Schweiz. Arch. Angew. Wiss. Tech.*, 1950, 18, 335-340; *Metal Abstr.*, 1951, 18, 675].—Two new hardness-testing machines for testing hard metals and very hard substances are described briefly. One is a static machine with a specially-shaped diamond indenter; the other is a microhardness tester in which the specimen is cut by a lightly-loaded diamond dust wheel rotating at high speed. R. B. CLARK.

Jim

GRODZINSKI, P.

Link Mechanisms in Modern
Kinematics

Instn mech.Engrs
Frep.

14pp.

1954

U.K.

P. Grodzinski, E. M'Ewen
A review of the investigations of linkages carried out
mainly in Germany and Russia. Possible future trends are
also discussed. (Bibl.57)

GRODZINSKI P.

✓ 466* Hardness Determination of Solid Bodies. Härteprüfung
fester Stoffe. 1. (German.) Grodzinski, Archiv für tech-
nische Messen, 1955, no. 230, Sept., p. 201-204.
Describes the principles and different methods of hardness
determination, static macro- and micro-hardness, scratch hard-
ness, and hardness scales. Graphs, tables. 82 ref.

JP
mst

Grodzinski P.

Low-Load Hardness, Its Elastic, Plastic, and Fracture Components. P. Grodzinski (*Metal*, 1955, 2, (13/14), 554-560).
The stress-strain relationship occurring during indentation hardness testing vary greatly according to the magnitude of the load; elastic and elasto-plastic stresses occur in micro and low-load testing, while plastic stresses occur in macro and macro-hardness testing. G. defines micro, low-load, and macro-hardness tests as being carried out, under loads of 1-200 g. (5-50 g.), 200 g.-10 kg. (200 g.-10 kg.), and >10 kg., resp., the values in brackets being those most usually encountered in practice. Plastic indentation by diamond pyramid indenters has led to the development of a new definition of hardness, which has as its basis the Meyer law, $A \propto F/d^2$. A method for determining the elastic component is described, and some of the results obtained thereby are discussed. During recent years many studies have been made of the nature of, and the hardness at which fracture of the diamond or of the immediate surroundings of the indentation takes place. The significance and the effects of such fractures on hardness determinations are described and illustrated; they are determined, to a large extent, by the type of indenter used—Vickers, Rockwell, Knoop, and double-load indenters being particularly employed. 25 refs. in text.

228 229

GRODZINSKI, W.

GRODZINSKI, W.

Fauna and flora in the Polish Beshchady Mountains.

P. 168 (Wierchy) Wol. 25, 1956, Kradow, Poland.

SO: MONTHLY INDEX OF EAST EUROPEAN ACCESSIONS (EEAI) LC. - VOL. 7, NO. 1, JAN. 1958

GRÓDZINCHI, W.

Studies of nature in the Beshchady Mountains.

P. 203 (Wierchy) Vol. 25, 1956, Krakow, Poland.

SO: MONTHLY INDEX OF EAST EUROPEAN ACCESSIONS (EEAI) LC. VOL. 7, NO. 1, JAN. 1958

GRÓDZINSKI, W.

GRÓDZINSKI, W.

Research on the successions of groups of small mammals in the forests of the Middle Beskids.

P. 205, Wierchy) Vol. 25, 1956, **Krakow**, Poland.

SO: MONTHLY INDEX OF EUROPEAN ACCESSIONS (EEAI) LC. VOL. 7, NO. 1, JAN. 1958

GRODZINSKI, W.

The succession of small mammal communities on an overgrown clearing and landslide in the Central Beskids of the western Carpathian Mountains. p. 83.

EKOLOGIA POLSKA. SERIA A. (Polska Akademia Nauk. Komitet Ekologiczny)
Warszawa, Poland. Vol. 7, no. 4, 1959.

Monthly List of East European Accessions (EEAI) LC, Vol. 9, no. 2, Feb. 1960

Uncl.

GRODZINSKI, W.

Metabolism rate and bioenergetics of small rodents from
the deciduous forest. Bul Ac Pol biol 9 no.12:493-499
'61.

1. Department of Animal Genetics and Organic Evolution,
Jagiellonian University, Cracow. Presented by T. Marchlewski.

*-

GRODZINSKI, Wladyslaw

Teodor Marchlewski, July 12, 1899 - January 27, 1962. Przegl
zoolog 6 no.4:253-258 '62.

1. Katedra Ewolucjonizmu i Genetyki Zwierzat, Uniwersytet
Jagiellobski, Krakow.

GRODZINSKI, Z.

1/2

Henr. K. Marcinkowski J., Grodziński Z. Skin-Drying of Moulding
Batches by Means of Infra-Red Rays

"Podawanie mas formierskich promieniami podczerwonymi". (Prace Inst. Odlewn. No. 4), Katowice, 1952, PWT, 18 pp., 30 figs.

Description of research carried out at the Foundry Institute over the drying of moulding batches by means of infra-red rays generated by Polish-made type PC 250-12 250-watt radiators. The experiments were based on recording the temperature in various layers of the moulding batch while it was being dried, and on determining, at various depths of the moulding batch, the remaining moisture content. The drying conditions were varied so as to be able to investigate the influence of the drying effect on all factors which play any role in the practical process of drying. It was found that the most economic results in drying mould with the aid of type PC-250-12 radiator will be obtained, if: 1) the mould enables the lamps to be located as close

(over)

21.

Notes K:

as possible in the surface; 2) the lumps are located as closely as possible one to the other; 3) the distance between the lumps and the surface which is being dried does not exceed 50 mm; 4) the angles between the axis of the lumps and the corresponding perpendiculars to the surface of the mould do not exceed 30°; 5) the supply of electric current is not susceptible to fall in voltage; 6) the moulding batch possesses good permeability and good tensile properties — both in the green and the dry state; 7) the correct moisture content of the moulding batch amounts to from 5 to 7 per cent H_2O ; 8) the moulding batch is coloured as dark as possible; 9) the depth of drying is fixed empirically for individual types of moulds and moulding batches, according to the quality of the castings and to the depth they occupy in the dried mould; 10) the optimum drying time is fixed so that it is, for individual drying conditions, in direct ratio to the required depth of drying; 11) the dried moulds are left to stand for a minimum time before being filled with metal.

Koss K., Gredziński Z., Marcinkowski J. Skin-Drying of Moulds by Means of Infra-Red Rays.

„Poduszanie form promieniami podczerwonymi”. (Prace Inst. Odlewn. No. 1), Stalingrad, 1953, PWT, 10 pp., 14 figs.

Description of laboratory and industrial experiments. Erection of industrial radiant heaters. Skin-drying of moulds. It has been found, by comparing skin-drying by infra-red ray heaters with the same process by ordinary coal fired drying ovens not arranged for recirculation of combustion products, that the use of radiant instead of box-type drying ovens slightly reduces the cost. The actual process of skin-drying by means of infra-red rays is more expensive than up-to-date air circulation and gas-fired box-type drying ovens, all the same the infra-red ray process has such advantages — notably the elimination of carriage of the moulds to the drying ovens — which are likely to prove economically decisive. An analysis of the advantages and disadvantages of this method shows that skin-drying of moulds by means of infra-red rays can be recommended in the following instances: 1) as a substitute for the dry-sand system of moulding large iron castings; 2) as a substitute for the dry-sand moulding of simple non-ferrous metal castings; 3) when drying patched-up parts of moulds already dried; 4) when working moulds from cement batches — as a means of accelerating the filling of the mould.

CRUZINSKI, Z.; HESS, K.

The use of inoculated cast iron for crane wheels. p. 103. (109-112 wanted)

(IRACE, Vol. 6, No. 3, 1956 (published 1957), Warszawa, Poland.)


SO: monthly List of East European Accessions (HEAL) Lc. Vol. 6, No. 10, October 1957. Uncl.

GRODZINSKI, Z.

Are migrating birds directed by the stars? p. 33

WSZECHSWIAT. (Polskie Towarzystwo Prsyrodnikow im. Kopernika)
Warszawa. No. 2, Feb. 1959
Poland/

Monthly List of East European Accessions Index (EEAI), IC, Vol. 8, No. 6, June 1959
Uncl.



PROCESSING AND PROPERTIES MODE	
BC	0
<p>Morphology of egg yolk under normal and experimental conditions. Z. GIMONIKI (Hull; Acad. Polonaise, 1955, 10, B, II, 317--355).—The reactions of yolk to f vapour, dioxan, and other vapour are described. The effect of vital staining is discussed with special reference to the lecithin content of yolk.</p> <p style="text-align: right;">W. F. R.</p>	
<p>ASD-55A METALLURGICAL LITERATURE CLASSIFICATION</p>	
<p>100000 MET ONLY 500</p>	<p>100000 MET ONLY 500</p>

Influence of the increase in the osmotic pressure upon
the white yolk spheres of the hen egg. *Z. für vergleichende
(Jagellonian Univ., Krakow). Bull. intern. Acad. polon.
Sci. Classe sci. math. nat. 1946, III, 87-98, cf. Bull.
Intern. Sci., Cracow, 1938. Drops of a suspension of yolk
of freshly laid hen egg in Pyralis's oil placed on cover
slips were mounted face down and allowed to evaporate. The
effect of desiccation on the white yolk spheres containing a
single drop of fat was then studied. Owing to the passage
of the osmotically free water through the semipermeable
membrane forming the wall of the sphere, the membrane
became viscous, as demonstrated by the formation of myo-
fibrin filaments and by adhesions of the spheres. Condensa-
tion of the protein moles, evidenced by turbidity, oc-
curred and was followed by a pptn. of the proteins and the
turning of the colloidal fluid into a granulous stiff mass.
The sphere diminished in size but maintained its shape.*

J. O. Holmes

CA

Digestion of the yolk of the hen egg. Z. Ciuplinski
(Jagellonian Univ., Krakow). *Skull. intern. acad. polon.*
111, 100 (1938) (in English).
Intercellular digestion of yolk was studied in tissue ob-
tained from the extravascular area of the yolk sac of 36-
44 hrs. incubated eggs and cultured in Tyrnole's skull.
Entodermal cells, with a plainly visible yolk sphere, were
examined every 3-5 hrs. The appearance of the yolk ele-
ments within the yolk sac during the entire time of incu-
bation, and the effect of the action of steapsin and pepsin on
the *in vitro* digestion of the yolk spheres were studied with
the aid of vital basic dyes and a polarizing microscope.
The yolk is an emulsion of the oil in water type, the con-
tinuous phase consisting of a protein soln. and the dis-
persed phase of glycerides drops. The yolk sphere con-
sists of both phases and of the superficial semipermeable
membrane. The latter is digested by lipase. Protease
acts on the continuous phase, aiding in its incorporation
into protoplasm. Lipase acts on the surface film of the

small fat drops, causing them to fuse into larger bodies,
and simultaneously transforms the glycerides into phos-
phatides. Various properties of yolk fat and yolk sphere
fat were established.

L. O. Holmes

BC

4 3-16

INFLUENCE OF TEMPERATURE ON RATE OF HEART IN EMBRYOS OF TELEOST
 (Fish. Ind. Acad. Polon. Sci., 1948, B.S.
 2. Grotowski).—The influence of temp. of 0°–45° on embry-
 onic heart was studied in several types of Teleost fish. Heart rate
 was found to increase with temp. and age of fish, the optimum
 temp. being between 17° and 20° depending on species. Changes in
 heart rate are predominantly controlled by myocardial factors. The
 heart is more resistant to unfavourable thermal conditions than
 other organs. M. K. Shood.

ASAC-55.6 METALLURGICAL LITERATURE CLASSIFICATION

Fat in the yolk of the sea trout *Salmo trutta*. J. Grodzinski (Jagellonian Univ., Krakow, Poland). *Bull. intern. acad. polon. sci., Classe sci. math. nat.* 1949 B:11. 50-78 (in English).—Yolk of the sea trout contains fat droplets which stain orange with Sudan III but do not stain with neutral red. During development to the free-swimming fish the fat droplets remain in the yolk and increase in size by fusion into larger drops. The position of fat drops changes with development. In the mature egg taken from the body cavity the fat is dispersed beneath the entire egg membrane except below the microvilli. After fertilization, at the time of formation of perivitellin fluid, most of the fat drops move toward the upper pole of the egg and collect beneath the germinal disk. Any remaining drops are pushed into the region of the embryo by the spreading yolk sac. The distribution of fat drops sets the position of the eggs in water and of the embryo within the egg membranes. The fat drops consist mostly of glycerides. The spherical shape of the fat drop is due to the presence of substances in the vitelline membrane — proteins and other N-containing compounds — which lower the vitelline fat interfacial tension. The drops flatten in physiological saline. Bifurcated needle-shaped crystals and centers of crystals appear within the fat drops after dilution of the yolk with Tyrode's fluid and strepsin. The fat drops then stain with neutral red. F. G. Shipley

CA

112

fat in the yolk of the sand lizard. Z. Gruzinski. Bull intern Acad polon sci, Classe sci math et nat Ser. B 10, 307-310, 1949 (in English). --Fat from the yolk sac of the lizard was investigated in a hanging drop, and studies made of its morphology and reactions to dyes (neutral red, trypan red, and Sudan III), osmotically active solns., and lipolytic ferment. Fat occurs in the yolk sac as drops of different sizes, shape, color, and sp. wt. Some drops consist probably of glycerides only, others are complexes of glycerides and proteins. The solely glyceride drops are digested by steapsin, and react only to Sudan III, but are resistant to osmotically active solns. and to neutral red. The complex drops stain with neutral red, and in hypotonic solns. vacuolize and break down into fat and protein substances. Under the influence of steapsin, the fat is segregated from

the protein component as minute granules. The sp. wt. of the complex fat drops is higher than that of the glyceride drops, and equals the sp. wt. of the red corpuscles of the embryo. Dorothy A. Meyer

GRODZINSKI, Zygmunt

Egg yolk of *Rhodeus amarus* Bl. Pol. morph. 5 no.1:13-26 1954.

1. Z Zakladu Anatomii Porownawczej im. H.Hoyera w Krakowie.
Kierownik: prof. dr Z.Grodzinski.

(FISH,

**Rhodeus amarus*, egg yolk)

(MOG YOLK,

*of *Rhodeus amarus*)

GRODZINSKI, Z.

Pulsation of different parts of a heart isolated from an embryo of the Salmo Trutta L. p. 65. FOLIA BIOLOGICA. (Panstwowe Wydawnictwo Naukowe), Warszawa. ol. 1, nos. 2-4, 1953. ol. 2, no. 3/4, 1954. DAFM Vol. 3, no. 1, 1955.

So. East European Accessions List. Vol. 5, no. 1, Jan. 1956

GRODZINSKI, Z.

/ 2478. Pulsation of different parts of the heart isolated from the sea-trout *Salmo trutta* L. embryos. Z. Grodzinski *Folia Biol.* Warsaw, 1955, 3, 65-82.—The hearts of the sea-trout embryos from 3 different developmental stages (I: 9-12 mm.; II: 15-18 mm.; III: 20-30 mm.) excised from the fish's body survive 2-3 days in Tyrode soln. adjusted for cold-blooded animals, with the addition of its own yolk. In such hearts the auricle was cut off from the ventricle or the connexion between them was only partially severed. The prep. were subjected to the influence of different temp. in special chambers. The current of water coming from the Hopper ultrathermostat maintained the desired temp. $\pm 0.1^\circ$. The pulsation was counted at 5°, 10°, 15°, 20°, 25°, 30°, and 34°. MD
B. YIMRY.

FOI b7D/ b7C and b7E and Partial P 76 (b) (7) (D) (b) (7) (C) (b) (7) (E) and b7E.

Doc. # 100: Ref. ZAMR-11 1.0.0.00 1978, 19229.

Author : Gr. Chisholm, AYC 113

151 :

1. Reaction of the Federal Court of Appeals of the Southern
Tribunal to Temperature Change.

Orig. Pub: *Encl. p14...* 1993-1995 (1995), 6, 15-3, 14-205.

Abstract: The isolated heart of the embryo of a salmon trout 12 - 29 cm long pulsated for 2 - 3 days in Tyrode's solution to which electrolytes of these same fish had been added, but the frequency of the contraction was less than in the embryo organism. With a temperature of 2 - 18 degrees the rate increased according to van't Hoff's law, at 18 - 30 degrees it did not change, with lethal temperatures (30 - 34 degrees) it decreased.

$$C_{-1}^2 : 1/2$$

FOUO/ Human and Animal Physiology. Blood Circulation. Heart.

T

Abs Jour: Ref Zhur-Biol., No 23, 1958, 93229.

and at 34 degrees the contractions ceased. The auricles of the isolated heart of younger embryos contracted more often than the ventricles (partial block). -- G.M. Cherkovich.

Card : 2/2

PROTEIN, L.

1453. Rhythmic rate of the heart of the sea-trout, *Salmo trutta* during embryonic development. Z. Grodzinski and A. Pigoń *Bull. Acad. polon. Sci.*, 1956, 4, 283-285 (Dept. of Comparat. Anat. Jagellonian Univ., Cracow, Poland).—Hearts excised from embryos of sea-trout survive for several hr. in Tyrode fluid; they react to changes in temp. by changes in the rhythm of pulsation. The older the embryo the quicker the rhythm. Hearts from younger embryos show peristaltic movements; those from older embryos systaltic contractions. Reduced wt. (and dry mass) of the heart grows rapidly during development from 0.55 µg. to 12.9 µg. Experimental methods are described and full data of measurements obtained are discussed.

P. HAAS

2

med

POLAND / General Biology. Individual Development.
Embryonic Development.

B

Abs Jour: Ref Zhur-Biol., No 23, 1958, 103289.

Author : Grodzinski, Zygmunt.

Inst : Not given.

Title : Provocation of Developmental Abnormalities of
Phylogenetic Significance.

Orig Pub: Kosmos, 1957, A6, No 1, 3-16.

Abstract: A review of works dealing with the influence of temperature on the formation of vertebrae in salmonoids and of the effect of insulin and other preparations in the formation of brachypodia, loss of the caudal portion of the skeleton and other abnormalities. The physical factors mentioned effect the development of the skeleton. This research has

Card 1/2

9

POLAND / General Biology. Individual Development.
Embryonic Development.

B

Ab's Jour: Ref Zhur-Biol., No 23, 1958, 103289.

Abstract: clarified at what developmental periods and by what agents individuals can be produced which differ from their parents. The altered animals could be considered subspecies or even species if the new characteristics could be transmitted by heredity. The author believes that this possibility exists.
-- V. A. Kanzyuba.

Card 2/2

GRODELSKI, Z.

Thoughts on vacations; a review of the collective work Problemy ewolucjonizmu. t. 2. Myśl ewolucyjna a paleozoologia (Problems of Evolutionism. Vol. 2. Evolutionary Thoughts and Paleozoology).

P. 633. (KOSMOS. SERIA A: BIOLOGIA) (Warszawa, Poland) Vol. 6, no. 6, 1957

OO: Monthly Index of East European Accession (MEEA) Vol. 7, No. 5, 1958

GRODZINSKI, Z.

Marine Biological Laboratory in Woods Hole, Mass. P 101

WSZECHSWIAT. (Polskie Towarzystwo Przyrodnikow im. Kopernika) Warszawa, 1959
No. 4, Apr. 1959

Monthly List of East European Accessions (EEAI) LC. Vol. 4, no. 7, July 1959

Uncl.

GRODZINSKI, Z.

The development of the lymph heart sea trout *Salmo trutta* L. Bul
Ac Pol biol 7 no.8:305-311 '59. (REAI 9:6)

1. Hoyer Department of Comparative Anatomy, Jagellonian University,
Krakow.
(Lymph) (Salmo trutta)

GRODZINS'KIY, A.M.; SITNIK, K.M.

"Study of growth promoting substances. Results and problems in research on growth promoting substances." H.Söding. Reviewed by A.M.Hrodzins'kyi, K.M.Sytnyk. Ukr.bot.zhur.13 no.2:115-118 '56.
(Growth promoting substances) (MIRA 9:9)

GRODZINSKIY, A.M., Cand Bio Sci --(diss) "Entrance and movement
of certain nutritive substances in plants under the ^{influence of soil} ~~effect~~ of pro-
^{cultivation} ~~cessing of soil~~ and fertilizers." Kiev, 1953. 15 pp (Acad Sci
USSR. Institute of Botany). 120 copies (SI, 33-52, 25)

SITNIK, K.M. [Sytnyk, K.M.]; GRODZINSKIY, A.M. [Grodzins'kyi, A.M.]

Effect of light on the absorption and translocation of phosphorus
in plants [with summary in English]. Ukr. bot. zhur. 15 no.2:29-35
'58. (MIRA 11:6)

1. Institut botaniki AN URSR, kafedra fiziologii roslin.
(Plants, Effect of light on) (Phosphorus) (Plants--Assimilation)

GRODZINSKIY, A.M. [Hrodzins'kyi, A.M.]; GRODZINSKIY, D.M. [Hrodzins'kyi, D.M.]

Relation between phosphorus and calcium uptake of corn plants and
the conditions of aeration. Ukr.bot.zhur. 15 no.4:3-10
'58. (MIRA 12:5)

1. Institut botaniki AN USSR i Ukrainskiy nauchno-issledovatel'-
skiy institut fiziologii rasteniy.
(Plants--Assimilation) (Plants--Respiration)
(Corn (Maize))

GRODZINSKIY, A.M. [Hrodzins'kyi, A.M.]; SYTHNIK, K.M. [Sytnyk, K.M.]

Recent data on the effect of light on nutrient uptake by
plants. Ukr.bot.zhur. 16 no.4:13-18 '59. (MIRA 12:11)

1. Institut botaniki AN USSR, otdel fiziologii rasteniy.
(Plants, Effect of light on)
(Plants, Motion of fluid in)

GRODZINSKIY, D.M. [Hrodzins'kyi, D.M.]; GRODZINSKIY, A.M. [Hrodzins'kyi, A.M.]

Methods of studying growth substances in plants; survey of literature. Ukr.bot.zhur. 16 no.4:51-66 '59.(MIRA 12:11)

1. Ukrainskiy nauchno-issledovatel'skiy institut fiziologii rasteniy Ukrainskoy akademii sel'skokhozyaystvennykh nauk i Institut botaniki AN USSR.
(Hormones (Plants))